

國立中山大學、國立成功大學合辦94學年度基礎學科
微積分競試試題與詳解

1. 試求 $\lim_{x \rightarrow \infty} \left(\frac{x^2-4}{x^2-1}\right)^{x^2+1}$.

Ans: Consider $\ln\left(\left(\frac{x^2-4}{x^2-1}\right)^{x^2+1}\right) = (x^2+1)(\ln(x^2-4) - \ln(x^2-1))$

$$\begin{aligned} \lim_{x \rightarrow \infty} (x^2+1)(\ln(x^2-4) - \ln(x^2-1)) &= \lim_{x \rightarrow \infty} \frac{\ln(x^2-4) - \ln(x^2-1)}{\frac{1}{x^2+1}} \\ \lim_{x \rightarrow \infty} \frac{\frac{2x}{x^2-4} - \frac{2x}{x^2-1}}{\frac{-2x}{(x^2+1)^2}} &= \lim_{x \rightarrow \infty} \frac{(x^2+1)^2(x^2-1-x^2+4)}{-(x^2-4)(x^2-1)} = -3 \end{aligned}$$

Since \ln is a continuous function, $\lim_{x \rightarrow a} \ln(x) = \ln(\lim_{x \rightarrow a} x)$, and by the L'Hospital's rule $\lim_{x \rightarrow \infty} \left(\frac{x^2-4}{x^2-1}\right)^{x^2+1} = e^{-3}$.

2. 試求 (a) $\int_0^1 x^5 e^{x^3} dx$.

$$(b) \int_0^{\frac{\pi}{4}} \frac{d\theta}{(\tan^2 \theta + 4 \tan \theta + 3) \cos^2 \theta}$$

$$(c) \int \frac{8x^2+4x-11}{(x+3)(x-1)^2} dx$$

$$\begin{aligned} \text{Ans: (a) Let } u = x^3. \text{ Then } \int_0^1 x^5 e^{x^3} dx &= \int_0^1 u \cdot \frac{1}{3} \cdot e^u du \\ &= \frac{1}{3} \left[ue^u \Big|_0^1 - \int_0^1 e^u du \right] = \frac{1}{3} \end{aligned}$$

$$(b) \text{ Let } u = \tan \theta \Rightarrow du = \sec^2 \theta d\theta$$

$$\begin{aligned} \text{原式} &= \int_0^1 \frac{du}{u^2+4u+3} = \int_0^1 \frac{1}{2} \left[\frac{1}{u+1} - \frac{1}{u+3} \right] du = \frac{1}{2} [\ln(u+1) - \ln(u+3)] \Big|_0^1 \\ &= \frac{1}{2} [\ln 3 - \ln 2] \end{aligned}$$

$$(c) \frac{8x^2+4x-11}{(x+3)(x-1)^2} = \frac{A}{x+3} + \frac{B}{x-1} + \frac{C}{(x-1)^2} \Rightarrow A = \frac{49}{16}, B = \frac{79}{16}, C = \frac{1}{4}$$

$$\text{原式} = \int \frac{\frac{49}{16}}{x+3} + \frac{\frac{79}{16}}{x-1} + \frac{\frac{1}{4}}{(x-1)^2} dx = \frac{49}{16} \ln|x+3| + \frac{79}{16} \ln|x-1| - \frac{1}{4} \frac{1}{x-1} + C$$

3. 試求 (a) $\int_{\frac{1}{2}}^1 \frac{1}{\sqrt{2t-t^2}} dt$ (b) $\lim_{x \rightarrow 1} \frac{1}{x-1} \left(\int_{\frac{1}{2}}^x \frac{1}{\sqrt{2t-t^2}} dt - \frac{\pi}{6} \right)$

Ans: (a) Letting $1-t = \sin \theta$, we have $dt = -\cos \theta d\theta$ and for $\frac{1}{2} \leq t \leq 1$, $0 \leq \sin \theta \leq \frac{1}{2} \Rightarrow \cos \theta \geq 0$ and $\sqrt{1-(1-t)^2} = \cos \theta$.

$$\text{Thus, } \int_{\frac{1}{2}}^1 \frac{1}{\sqrt{1-(1-t)^2}} dt = \int_{\frac{\pi}{6}}^0 \frac{-\cos \theta d\theta}{\cos \theta} = -\theta \Big|_{\frac{\pi}{6}}^0 = \frac{\pi}{6}$$

(b) Let $f(x) = \int_{\frac{1}{2}}^x \frac{1}{\sqrt{2t-t^2}} dt$. Then $\lim_{x \rightarrow 1} \frac{1}{x-1} \left(\int_{\frac{1}{2}}^x \frac{1}{\sqrt{2t-t^2}} dt - \frac{\pi}{6} \right) = \lim_{x \rightarrow 1} \frac{f(x)-f(1)}{x-1}$.

Since $\sqrt{2t-t^2} > 0$ for $0 < t < 2$, $f(x)$ is differentiable at $x = 1$, and the

Fundamental Theorem of Calculus implies that

$$\lim_{x \rightarrow 1} \frac{f(x) - f(1)}{x - 1} = f'(1) = \left. \frac{1}{\sqrt{2t-t^2}} \right|_{t=1} = 1$$

4. 請判定瑕積分 $\int_0^\infty e^{x-2e^x} dx$ 是否存在？如果此瑕積分存在，試求其值。


$$\begin{aligned} \text{Ans: Consider } & \int_0^t \frac{e^x}{e^{2e^x}} dx. \quad (\text{Let } u = e^x) \\ & = \int_1^{e^t} \frac{du}{e^{2u}} = \int_1^{e^t} -\frac{1}{2}(-2e^{-2u}) du \\ & = -\frac{1}{2}e^{-2u} \Big|_1^{e^t} = -\frac{1}{2}(e^{-2e^t} - e^{-2}) \end{aligned}$$

$$\lim_{t \rightarrow \infty} -\frac{1}{2}(e^{-2e^t} - e^{-2}) = \lim_{t \rightarrow \infty} \frac{-1}{2} \left(\frac{1}{e^{2e^t}} - \frac{1}{e^2} \right) = \frac{1}{2e^2}$$

5. 令 $R = \{(x, y) \in \mathbb{R}^2 | 1 \leq x \leq 4, 1 < y \leq 1 + x^{\frac{1}{2}}\}$. 試求：

- (a) R 對 $x = 1$ 旋轉所得之旋轉體體積.
- (b) R 對 $x = 0$ 旋轉所得之旋轉體體積.

Ans:



$$\begin{aligned} \text{(a) } Vol &= \int_1^4 2\pi(x-1)[1 + \sqrt{x} - 1]dx \\ &= \int_1^4 2\pi(x^{\frac{3}{2}} - x^{\frac{1}{2}})dx \end{aligned}$$

$$= 2\pi \left(\frac{2}{5}x^{\frac{5}{2}} - \frac{2}{3}x^{\frac{3}{2}} \right) \Big|_1^4$$

$$= 2\pi \left(\frac{62}{5} - \frac{14}{3} \right)$$

另解: $Vol = \pi \cdot 3^2 + \int_2^3 \pi \{3^2 - [(y-1)^2 - 1]^2\} dy$
 $= \pi \cdot 3^2 \cdot 2 - \int_2^3 \pi [(y-1)^2 - 1]^2 dy$

(b) $Vol = \int_1^4 2\pi x [1 + \sqrt{x} - 1] dx$
 $= \int_1^4 2\pi x^{\frac{3}{2}} dx$
 $= \frac{4\pi}{5} x^{\frac{5}{2}} \Big|_1^4 = \frac{124\pi}{5}$

另解: $Vol = \pi(4^2 - 1^2) + \int_2^3 \pi [4^2 - (y-1)^4] dy$
 $= \pi \cdot 4^2 \cdot 2 - \pi \cdot 1^2 \cdot 1 - \int_2^3 \pi [(y-1)^2]^2 dy$

6. 考慮曲線 $y = x^{\frac{1}{2}}$ 對 $y = 0$ 旋轉. 試求當 $1 \leq x \leq 4$ 時所得之旋轉體表面積.

Ans: $Area = \int_1^4 2\pi \cdot x^{\frac{1}{2}} \cdot \sqrt{1 + (\frac{1}{2}x^{-\frac{1}{2}})^2} dx = \int_1^4 2\pi \cdot x^{\frac{1}{2}} \cdot \sqrt{1 + \frac{1}{4}x^{-1}} dx$
 $= 2\pi \int_1^4 \sqrt{x + \frac{1}{4}} dx = 2\pi (x + \frac{1}{4})^{\frac{3}{2}} \cdot \frac{2}{3} \Big|_1^4$
 $= \frac{4\pi}{3} (\frac{17^{\frac{3}{2}}}{8} - \frac{5^{\frac{3}{2}}}{8}) = \frac{\pi}{6} (17^{\frac{3}{2}} - 5^{\frac{3}{2}})$

7. 設 $f(x) = \sqrt{x^3 + x + 6}$, 求 $(f^{-1})'(4)$, 也就是求 f 的反函數在 4 的導數.

Ans: $4 = \sqrt{x^3 + x + 6} \Rightarrow x = 2$ (另一根 $x^2 + 2x + 5$ 虛根)
 $\therefore (f^{-1})'(4) = \frac{1}{f'(2)} = \frac{1}{\frac{13}{8}} = \frac{8}{13}$

8. 設 $f(x) = \begin{cases} 3x + x^2 \sin \frac{1}{x} & \text{當 } x \neq 0 \\ 0 & \text{當 } x = 0 \end{cases}$

請判定 $f'(x)$ 存在的位置. 當 $f'(x)$ 存在時, 試求其公式.

Ans: When $x \neq 0$, $f'(x) = 3 + 2x \sin \frac{1}{x} + x^2 \cos \frac{1}{x} \cdot (-\frac{1}{x^2})$
 $= 3 + 2x \sin \frac{1}{x} - \cos \frac{1}{x}$

At $x = 0$, $\lim_{x \rightarrow 0} \frac{f(x) - f(0)}{x - 0} = \lim_{x \rightarrow 0} \frac{3x + x^2 \sin \frac{1}{x}}{x}$
 $= \lim_{x \rightarrow 0} (3 + x \sin \frac{1}{x})$

And, since $|\sin \frac{1}{x}| \leq 1$, $0 \leq |x \sin \frac{1}{x}| \leq |x|$

$\Rightarrow \lim_{x \rightarrow 0} 0 \leq \lim_{x \rightarrow 0} |x \sin \frac{1}{x}| \leq \lim_{x \rightarrow 0} |x| = 0$

$\Rightarrow \lim_{x \rightarrow 0} |x \sin \frac{1}{x}| = 0$

$\Rightarrow \lim_{x \rightarrow 0} x \sin \frac{1}{x} = 0$

$\Rightarrow f'(0) = 3$

9. 令 $f(x) = \frac{x^3}{x^2 - 1}$. 請討論 $f(x)$ 遞增, 遞減, 凸凹, 漸近線, 極值之位置, 並畫出 $y = f(x)$ 的圖.

Ans: 漢近線; $x = 1, x = -1, y = x$.

local maximum: $(-\sqrt{3}, -\frac{3\sqrt{3}}{2})$.

local minimum: $(\sqrt{3}, \frac{3\sqrt{3}}{2})$.

point of reflection: $(0, 0)$.

圖形:

